DEFINITION

APPLYING PLANT RESIDUES OR OTHER SUITABLE MATERIALS, PRODUCED ON THE SITE IF POSSIBLE, TO THE SOIL SURFACE.

PURPOSE:

- TO REDUCE RUNOFF AND EROSION
- TO CONSERVE MOISTURE
- TO PREVENT SURFACE COMPACTION OR CRUSTING
- TO CONTROL UNDESIRABLE VEGETATION
- TO MODIFY SOIL TEMPERATURE
- TO INCREASE BIOLOGICAL ACTIVITY IN THE SOIL

REQUIREMENT FOR REGULATORY COMPLIANCE

MULCH OR TEMPORARY GRASSING SHALL BE APPLIED TO ALL EXPOSED AREAS WITHIN 14 DAYS OF DISTURBANCE. MULCH CAN BE USED AS A SINGULAR EROSION CONTROL DEVICE FOR UP TO SIX MONTHS, BUT IT SHALL BE APPLIED AT THE APPROPRIATE DEPTH, DEPENDING ON THE MATERIAL USED, AND/OVERLAID WITH A CONTINUOUS 90% COVER OR GREATER OF THE SOIL SURFACE.

MAINTENANCE SHALL BE REQUIRED TO MAINTAIN APPROPRIATE DEPTH AND 90% COVER. TEMPORARY VEGETATION MAY BE EMPLOYED INSTEAD OF MULCH IF THE AREA WILL REMAIN UNDISTURBED FOR LESS THAN SIX MONTHS.

IF ANY AREA WILL REMAIN UNDISTURBED FOR GREATER THAN SIX MONTHS, PERMANENT VEGETATIVE TECHNIQUES SHALL BE EMPLOYED. REFER TO DS2 –DISTURBED AREA STABILIZATION (WITH TEMPORARY SEEDING), DS3 – DISTURBED AREA STABILIZATION (WITH PERMANENT SEEDING), AND DS4 – DISTURBED AREA STABILIZATION (WITH SODDING).

SPECIFICATIONS

MULCHING WITHOUT SEEDING

THIS STANDARD APPLIES TO GRADED OR CLEARED AREAS WHERE SEEDINGS MAY NOT HAVE A SUITABLE GROWING SEASON TO PRODUCE AN EROSION RESISTANT COVER, BUT CAN BE STABILIZED WITH A MULCH COVER.

SITE PREPARATION

1. GRADE TO PERMIT THE USE OF EQUIPMENT FOR APPLYING AND ANCHORING MULCH.
2. INSTALL NEEDED EROSION CONTROL MEASURES AS REQUIRED SUCH AS DIKES, DIVERSIONS, SEDIMENT BARRIERS AND SEDIMENT BARRIERS.
3. LOOSEN COMPACT SOIL TO A MINIMUM DEPTH OF 3 INCHES.

MULCHING MATERIALS

SELECT ONE OF THE FOLLOWING MATERIALS AND APPLY AT THE DEPTH INDICATED:

1. DRY STRAW OR HAY SHALL BE APPLIED AT A DEPTH OF 2 TO 4 INCHES PROVIDING COMPLETE SOIL COVERAGE. ONE ADVANTAGE OF THIS MATERIAL IS EASY APPLICATION.
2. WOOD WASTE (CHIPS, SAWDUST OR BARK) SHALL BE APPLIED AT A DEPTH OF 2 TO 3 INCHES. ORGANIC MATERIAL FROM THE CLEARING STAGE OF DEVELOPMENT SHOULD REMAIN ON SITE, BE CHIPPED, AND APPLIED AS MULCH. THIS METHOD OF MULCHING CAN GREATLY REDUCE EROSION CONTROL COSTS.
3. POLYETHYLENE FILM SHALL BE SECURED OVER BANKS OR STOCKPILED SOIL MATERIAL FOR TEMPORARY PROTECTION. THIS MATERIAL CAN BE SALVAGED AND RE-USED.

APPLYING MULCH

WHEN MULCH IS USED WITHOUT SEEDING, MULCH SHALL BE APPLIED TO PROVIDE FULL COVERAGE OF THE EXPOSED AREA.

1. DRY STRAW OR HAY MULCH AND WOOD CHIPS SHALL BE APPLIED UNIFORMLY BY HAND OR BY MECHANICAL EQUIPMENT.
2. IF THE AREA WILL EVENTUALLY BE COVERED WITH PERENNIAL VEGETATION, 20–30 POUNDS OF NITROGEN PER ACRE IN ADDITION TO THE NORMAL AMOUNT SHALL BE APPLIED TO OFFSET THE UPTAKE OF NITROGEN CAUSED BY THE DECOMPOSITION OF THE ORGANIC MULCHES.
3. APPLY POLYETHYLENE FILM ON EXPOSED AREAS.

ANCHORING MULCH

1. STRAW OR HAY MULCH CAN BE PRESSED INTO THE SOIL WITH A DISK HARROW WITH THE DISK SET STRAIGHT OR WITH A SPECIAL "PACKER" DISK. DISKS MAY BE SMOOTH OR SERRATED AND SHOULD BE 20 INCHES OR MORE IN DIAMETER AND 8 TO 12 INCHES APART. THE EDGES OF THE DISK SHOULD BE DULL ENOUGH NOT TO CUT THE MULCH BUT TO PRESS IT INTO THE SOIL LEAVING MUCH OF IT IN AN ERECT POSITION. STRAW OR HAY MULCH SHALL BE ANCHORED IMMEDIATELY AFTER APPLICATION.
2. STRAW OR HAY MULCH SPREAD WITH SPECIAL BLOWER-TYPE EQUIPMENT MAY BE ANCHORED BY TACKIFIERS, BINDERS AND HYDRAULIC MULCH WITH TACKIFIER SPECIFICALLY DESIGNED TO TACKIFY STRAW CAN BE SUBSTITUTED FOR EMULSIFIED ASPHALT. PLEASE REFER TO SPECIFICATION TAC–TACKIFIERS. PLASTIC MESH OR NETTING WITH MESH NO LARGER THAN ONE INCH BY ONE INCH SHALL BE INSTALLED ACCORDING TO MANUFACTURER'S SPECIFICATIONS.
3. NETTING OF THE APPROPRIATE SIZE SHALL BE USED TO ANCHOR WOOD WASTE. OPENINGS OF THE NETTING SHALL NOT BE LARGER THAN THE AVERAGE SIZE OF THE WOOD WASTE CHIPS.
4. POLYETHYLENE FILM SHALL BE ANCHOR TRENCHED AT THE TOP AS WELL AS INCREMENTALLY AS NECESSARY.

Ds1 DISTURBED AREA STABILIZATION (WITH MULCHING ONLY)
LOCATION IN MANUAL : 6-27

DEFINITION

THE ESTABLISHMENT OF TEMPORARY VEGETATIVE COVER WITH FAST GROWING SEEDINGS FOR SEASONAL PROTECTION ON DISTURBED OR DENUDED AREAS.

PURPOSE:

- TO REDUCE RUNOFF AND SEDIMENT DAMAGE OF DOWN STREAM RESOURCES
- TO PROTECT THE SOIL SURFACE FROM EROSION
- TO IMPROVE WILDLIFE HABITAT
- TO IMPROVE AESTHETICS
- TO IMPROVE TILTH, INFILTRATION AND AERATION AS WELL AS ORGANIC MATTER FOR PERMANENT PLANTINGS

REQUIREMENT FOR REGULATORY COMPLIANCE

MULCH OR TEMPORARY GRASSING SHALL BE APPLIED TO ALL EXPOSED AREAS WITHIN 14 DAYS OF DISTURBANCE. TEMPORARY GRASSING, INSTEAD OF MULCH, CAN BE APPLIED TO ROUGH GRADED AREAS THAT WILL BE EXPOSED FOR LESS THAN SIX MONTHS. IF AN AREA IS EXPECTED TO BE UNDISTURBED FOR LONGER THAN SIX MONTHS, PERMANENT PERENNIAL VEGETATION SHALL BE USED. IF OPTIMUM PLANTING CONDITIONS FOR TEMPORARY GRASSING IS LACKING, MULCH CAN BE USED AS A SINGULAR EROSION CONTROL DEVICE FOR UP TO SIX MONTHS BUT IT SHALL BE APPLIED AT THE APPROPRIATE DEPTH, ANCHORED, AND HAVE A CONTINUOUS 90% COVER OR GREATER OF THE SOIL SURFACE. REFER TO SPECIFICATION DS1–DISTURBED AREA STABILIZATION (WITH TEMPORARY SEEDING).

CONDITIONS

TEMPORARY VEGETATIVE MEASURES SHOULD BE COORDINATED WITH PERMANENT MEASURES TO ASSURE ECONOMICAL AND EFFECTIVE STABILIZATION. MOST TYPES OF TEMPORARY VEGETATION ARE IDEAL TO USE AS COMPANION CROPS UNTIL THE PERMANENT VEGETATION IS ESTABLISHED. NOTE: SOME SPECIES OF TEMPORARY VEGETATION ARE NOT APPROPRIATE FOR COMPANION CROP PLANTINGS BECAUSE OF THEIR POTENTIAL TO OUT–COMPETE THE DESIRED SPECIES (E.G. ANNUAL HYGRASS). CONTACT NRCS OR THE LOCAL SWCD FOR MORE INFORMATION.

SPECIFICATIONS

GRADING AND SHAPING

EXCESSIVE WATER RUN–OFF SHALL BE REDUCED BY PROPERLY DESIGNED AND INSTALLED EROSION CONTROL PRACTICES SUCH AS CLOSED DRAINS, DITCHES, DIKES, DIVERSIONS, SEDIMENT BARRIERS AND OTHERS.

NO SHAPING OR GRADING IS REQUIRED IF SLOPES CAN BE STABILIZED BY HAND–SEEDED VEGETATION OR IF HYDRAULIC SEEDING EQUIPMENT IS TO BE USED.

SEEDBED PREPARATION

WHEN A HYDRAULIC SEEDER IS USED, SEEDBED PREPARATION IS NOT REQUIRED. WHEN USING CONVENTIONAL OR HAND–SEEDING, SEEDBED PREPARATION IS NOT REQUIRED IF THE SOIL MATERIAL IS LOOSE AND NOT SEALED BY RAINFALL.

WHEN SOIL HAS BEEN SEALED BY RAINFALL OR CONSISTS OF SMOOTH CUT SLOPES, THE SOIL SHALL BE PITTED, TRENCHED OR OTHERWISE SCARIFIED TO PROVIDE A PLACE FOR SEED TO LODGE AND GERMINATE.

LIME AND FERTILIZER

AGRICULTURAL LIME IS REQUIRED UNLESS SOIL TESTS INDICATE OTHERWISE. APPLY AGRICULTURAL LIME AT A RATE DETERMINED BY SOIL TEST FOR PH. QUICK ACTING LIME SHOULD BE INCORPORATED TO MODIFY PH DURING THE GERMINATION PERIOD. 80 STIMULANTS SHOULD ALSO BE CONSIDERED WHEN THERE IS LESS THAN 3% ORGANIC MATTER IN THE SOIL. GRADED AREAS REQUIRE LIME APPLICATION. SOILS MUST BE TESTED TO DETERMINE REQUIRED AMOUNTS OF FERTILIZER AND AMENDMENTS. FERTILIZER SHOULD BE APPLIED BEFORE LAND PREPARATION AND INCORPORATED WITH A DISC, RIPPER, OR CHISEL. ON SLOPES TOO STEEP FOR, OR INACCESSIBLE TO EQUIPMENT, FERTILIZER SHALL BE HYDRAULICALLY APPLIED, PREFERABLY IN THE FIRST PASS WITH SEED AND SOME HYDRAULIC MULCH, THEN TOPPED WITH THE REMAINING REQUIRED APPLICATION RATE.

SEEDING

SELECT A GRASS OR GRASS–LEGUME MIXTURE SUITABLE TO THE AREA AND SEASON OF THE YEAR. SEED SHALL BE APPLIED UNIFORMLY BY HAND, CYCLONE SEEDER, DRILL, CULTI–PACKER–SEEDER, OR HYDRAULIC SEEDER (SLURRY INCLUDING SEED AND FERTILIZER). DRILL OR CULTIPACKER SEEDING SHALL NORMALLY PLACE SEED ONE–QUARTER TO ONE–HALF INCH DEEP. MAXIMUM DEPTH OF PLANTING IS TEN TIMES THE SEED DIAMETER, SMALL SEEDS BE "RAKED" LIGHTLY TO COVER SEED WITH SOIL IF SEEDED BY HAND. SEE TABLE 6–4.1

MULCHING

TEMPORARY VEGETATION CAN, IN MOST CASES, BE ESTABLISHED WITHOUT THE USE OF MULCH, PROVIDED THERE IS LITTLE TO NO EROSION POTENTIAL. HOWEVER, THE USE OF MULCH CAN GREATLY ACCELERATE AND ENHANCE GERMINATION AND VEGETATIVE ESTABLISHMENT. MULCH WITHOUT SEEDING SHOULD BE CONSIDERED FOR SHORT TERM PROTECTION. REFER TO DS1 – DISTURBED AREA STABILIZATION (WITH MULCHING ONLY).

IRRIGATION

DURING TIMES OF DROUGHT, WATER SHALL BE APPLIED AT A RATE NOT CAUSING RUNOFF AND EROSION. THE SOIL SHALL BE THOROUGHLY WETTED TO A DEPTH THAT WILL INSURE GERMINATION OF THE SEED. SUBSEQUENT APPLICATIONS SHOULD BE MADE WHEN NEEDED.

Ds2 DISTURBED AREA STABILIZATION (WITH TEMPORARY SEEDING)
LOCATION IN MANUAL : 6-29

TABLE 6–4.1
PLANTS, PLANTING RATES, AND PLANTING DATES FOR TEMPORARY COVER OR COMPANION CROPS

SPECIES	BROADCAST per ** Acre	BROADCAST PLS * per 1000 SF	RESOURCE	REMARKS
LESPEDEZA ANNUAL (LESPEDEZA STRIATA) ALONE ..... IN MIXTURES	40 lb. 10 lb.	0.9 lb. 0.2 lb.	MOUNTAINS SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	200,000 SEED PER POUND MAY VOLUNTEER FOR SEVERAL YEARS USE INOCULANT (L)
LOVEGRASS, WEEPING (ERAGROSTIS CURVULA) ALONE ..... IN MIXTURES	4 lbs. 2 lbs.	0.1 lb. 0.05 lb.	MOUNTAINS SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	1,300,000 SEED PER POUND MAY LAST FOR SEVERAL YEARS MIX W/ SERICEA LESPEDEZA
MILLET, BROWNTOP (PANICUM FASCICULATUM) ALONE ..... IN MIXTURES	40 lbs. 10 lbs.	0.9 lb. 0.2 lb.	MOUNTAINS SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	137,000 SEED PER POUND QUICK GROWING COVER WILL PROVIDE TOO MUCH COMPETITION IN MIXTURES IF SEEDED AT HIGH RATES
RYE (SECALE CEREALE) ALONE ..... IN MIXTURES	3 bu. (168 lbs.) 1/2 bu. (28 lbs.)	3.9 lb. 0.6 lb.	MOUNTAINS SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	18,000 SEED PER POUND STOCK COVER CROSSES TOLERANT AND WINTER-HARDY
RYEGRASS, ANNUAL (LOLIUM TEMULENTUM) ALONE	40 lbs.	0.9 lb.	MOUNTAINS SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	227,000 SEED PER POUND DONOT COVER VERY COMPETITIVE AND IS NOT TO BE USED IN MIXTURES

BOLD LINES (▬) INDICATED OPTIMUM DATES, GREY LINES (▭) INDICATED PERMISSIBLE BUT MARGINAL DATES
* PLS – PURE LIVE SEED
** REDUCE RATES BY 50% WHEN DRILLED.

Ds2 DISTURBED AREA STABILIZATION (WITH TEMPORARY SEEDING)
LOCATION IN MANUAL : 6-31

PLANTING AND SEEDING SPECIFICATIONS

PLANT SELECTION

REFER TO TABLES 6–4.1, 6–5.2, 6–5.3 AND 6–5.4 FOR APPROVED SPECIES. SPECIES NOT LISTED MAY BE APPROVED BY THE STATE RESOURCE CONSERVATIONIST OF THE NATURAL RESOURCES CONSERVATION SERVICE BEFORE THEY ARE USED.

PLANTS SHALL BE SELECTED ON THE BASIS OF SPECIES CHARACTERISTICS, SITE AND SOIL CONDITIONS, PLANNED USE AND MAINTENANCE OF THE AREA; TIME OF YEAR OF PLANTING, METHOD OF PLANTING; AND THE NEEDS AND DESIRES OF THE LAND USER.

SOME PERENNIAL SPECIES ARE EASILY ESTABLISHED AND CAN BE PLANTED ALONE. EXAMPLES OF THESE ARE COMMON BERMUDA, TALL FESCUE, AND WEEPING LOVEGRASS.

OTHER PERENNIALS, SUCH AS BAHIA GRASS AND SERICEA LESPEDEZA, ARE SLOW TO BECOME ESTABLISHED AND SHOULD BE PLANTED WITH ANOTHER PERENNIAL SPECIES. THE ADDITIONAL SPECIES WILL PROVIDE QUICK COVER AND AMPLE SOIL PROTECTION UNTIL THE TARGET PERENNIAL SPECIES BECOME ESTABLISHED. FOR EXAMPLE, COMMON SEEDING COMBINATIONS ARE 1) WEEPING LOVEGRASS WITH SERICEA LESPEDEZA (SCARIFIED) AND 2) TALL FESCUE WITH SERICEA LESPEDEZA (UNSCARIFIED).

PLANT SELECTION MAY ALSO INCLUDE ANNUAL COMPANION CROPS. ANNUAL COMPANION CROPS SHOULD BE USED ONLY WHEN THE PERENNIAL SPECIES ARE NOT PLANTED DURING THEIR OPTIMUM PLANTING PERIOD. A COMMON COMPANION CROP MIXTURE IS MILLET WITH COMMON BERMUDA IN MID–SUMMER. CARE SHOULD BE TAKEN IN SELECTING A COMPANION CROP SPECIES AND SEEDING RATES BECAUSE ANNUAL CROPS WILL COMPETE WITH PERENNIAL SPECIES FOR WATER, NUTRIENTS, AND GROWING SPACE. A HIGH SEEDING RATE OF A COMMON CROP MAY PREVENT THE ESTABLISHMENT OF PERENNIAL SPECIES.

RYEGRASS SHALL NOT BE USED IN ANY SEEDING MIXTURES CONTAINING PERENNIAL SPECIES DUE TO ITS ABILITY TO OUT–COMPETE DESIRED SPECIES CHOSEN FOR PERMANENT PERENNIAL COVER.

SEED QUALITY

THE TERM "PURE LIVE SEED" IS USED TO EXPRESS THE QUALITY OF SEED AND IS NOT SHOWN ON THE LABEL. PURE LIVE SEED, PLS, IS EXPRESSED AS A PERCENTAGE OF THE SEEDS THAT ARE PURE AND WILL GERMINATE. INFORMATION ON PERCENT GERMINATION AND PURITY CAN BE FOUND ON SEED TAGS. PLS IS DETERMINED BY MULTIPLYING THE PERCENT OF PURITY TIMES THE PERCENT OF GERMINATION; I.E.,

(PLS = % GERMINATION X % PURITY)

EXAMPLE:

COMMON BERMUDA SEED
70% GERMINATION, 80% PURITY

PLS = 70% GERMINATION X 80% PURITY

PLS = 56%

THE PERCENT OF PLS HELPS YOU DETERMINE THE AMOUNT OF SEED YOU NEED. IF THE SEEDING RATE IS TO POUNDS PLS AND THE BULK SEED IS 56 % PLS, THE BULK SEEDING RATE IS:

10 LBS. PLS/ACRE ÷ 17.9 LBS/ACRE
56% PLS

YOU WOULD NEED TO PLANT 17.8 LBS/ACRE TO PROVIDE 10 LBS/ACRE OF PURE LIVE SEED.

SEEDBED PREPARATION

SEEDBED PREPARATION MAY NOT BE REQUIRED WHERE HYDRAULIC SEEDING AND FERTILIZING EQUIPMENT IS TO BE USED (BUT IS STRONGLY RECOMMENDED FOR ANY SEEDING PROCESS, WHEN POSSIBLE), WHEN CONVENTIONAL SEEDING IS TO BE USED, SEEDBED PREPARATION WILL BE DONE AS FOLLOWS:

BROADCAST PLANTINGS

1. *TILLAGE*, AT A MINIMUM, SHALL ADEQUATELY 6–90 LOOSEN THE SOIL TO A DEPTH OF 4 TO 6 INCHES. ALLEVIATE SOIL COMPACTION, INCORPORATE LIME AND FERTILIZER; SMOOTH AND FIRM THE SOIL; ALLOW FOR THE PROPER PLACEMENT OF SEED, SPROGS, OR PLANTS; AND ALLOW FOR THE INCORPORATING OF STRAW OR HAY MULCH IF A DISK IS TO BE USED.
2. *TILLAGE* MAY BE DONE WITH ANY SUITABLE EQUIPMENT.
3. *TILLAGE* SHOULD BE DONE ON THE CONTOUR WHERE FEASIBLE.
4. ON SLOPES TOO STEEP FOR THE SAFE OPERATION OF TILLAGE EQUIPMENT, THE SOIL SURFACE SHALL BE PITTED OR TRENCHED ACROSS THE SLOPE WITH APPROPRIATE HAND TOOLS TO PROVIDE TWO PLACES FOR EACH SEED, SPACED 6 INCHES APART IN WHICH SEED MAY LODGE AND GERMINATE. HYDRAULIC SEEDING MAY ALSO BE USED.

INDIVIDUAL PLANTS

1. WHERE INDIVIDUAL PLANTS ARE TO BE SET, THE SOIL SHALL BE PREPARED BY EXCAVATING HOLES, OPENING FURROWS, OR DIBBLE PLANTING.
2. FOR NURSERY STOCK PLANTS, HOLES SHALL BE LARGE ENOUGH TO ACCOMMODATE ROOTS WITHOUT CROWDING.
3. WHERE PINE SEEDLINGS ARE TO BE PLANTED, SUBSOIL UNDER THE ROW 36 INCHES DEEP ON THE CONTOUR FOUR TO SIX MONTHS PRIOR TO PLANTING. SUBSOILING SHOULD BE DONE WHEN THE SOIL IS DRY, PREFERABLY IN AUGUST OR SEPTEMBER.

INNOCULANTS

ALL LEGUME SEED SHALL BE INOCULATED WITH APPROPRIATE NITROGEN–FIXING BACTERIA. THE INNOCULANT SHALL BE A PURE CULTURE PREPARED SPECIFICALLY FOR THE SEED SPECIES AND USED WITHIN THE DATES ON THE CONTAINER.

A MIXING MEDIUM RECOMMENDED BY THE MANUFACTURER SHALL BE USED TO BOND THE INNOCULANT TO THE SEED. FOR CONVENTIONAL SEEDING, USE TWICE THE AMOUNT OF INNOCULANT RECOMMENDED BY THE MANUFACTURER. FOR HYDRAULIC SEEDING, USE 4 TIMES THE AMOUNT OF INNOCULANT RECOMMENDED BY THE MANUFACTURER SHAKE WELL.

ALL INOCULATED SEED SHALL BE PROTECTED FROM THE SUN AND DRYING CONDITIONS. INOCULATED SEED SHALL REMAIN MOIST AND HYDROSEEDED WITHIN 24 HOURS AFTER MIXING.

SEEDING

HYDRAULIC SEEDING

MIX THE SEED (INNOCULATED IF NEEDED), FERTILIZER, AND WOOD CELLULOSE OR WOOD PULP FIBER MULCH WITH WATER AND APPLY IN A SLURRY UNIFORMLY OVER THE AREA TO BE TREATED. APPLY WITHIN ONE HOUR AFTER THE SLURRY IS MADE.

CONVENTIONAL SEEDING

SEEDING WILL BE DONE ON A FRESHLY PREPARED AND FIRMED SEEDBED. FOR BROADCAST PLANTING, USE A CULTI–PACKER–SEEDER, DRILL, ROTARY SEEDER, OTHER MECHANICAL SEEDER, OR HAND SEEDING TO DISTRIBUTE THE SEED UNIFORMLY OVER THE AREA TO BE TREATED. COVER THE SEED LIGHTLY WITH 1/8 TO 1/4 INCH OF SOIL FOR SMALL SEED AND 1/2 TO 1 INCH FOR LARGE SEED WHEN USING A CULTIPACKER OR OTHER SUITABLE EQUIPMENT.

Ds3 DISTURBED AREA STABILIZATION (PERMANENT VEGETATION)
LOCATION IN MANUAL : 6-37

MULCHING RATES AND SPECIFICATIONS

MULCH IS REQUIRED FOR ALL PERMANENT VEGETATION APPLICATIONS. MULCH APPLIED TO SEEDED AREAS SHALL ACHIEVE 75% TO 100% SOIL COVER. WHEN SELECTING A MULCH, DESIGN PROFESSIONALS SHOULD CONSIDER MULCH'S FUNCTIONAL LONGEVITY, EROSION CONTROL ENHANCEMENT, AND EROSION CONTROL REQUIREMENTS. SELECT THE MULCHING MATERIAL FROM THE FOLLOWING AND APPLY AS INDICATED:

1. DRY STRAW OR HAY MULCH OF GOOD QUALITY AND FREE OF WEED SEEDS CAN BE USED. STRAW SHALL BE APPLIED AT THE RATE OF 2 TONS PER ACRE. DRY HAY SHALL BE APPLIED AT A RATE OF 2 1/2 TONS PER ACRE.
2. WOOD CELLULOSE FIBER OR WOOD PULP FIBER SHALL BE USED IN HYDRAULIC SEEDING. IT SHALL BE APPLIED AT THE RATE OF 500 POUNDS PER ACRE. DRY STRAW OR DRY HAY MULCH SHALL BE APPLIED (AT THE RATE INDICATED ABOVE) AFTER HYDRAULIC SEEDING.
3. ONE THOUSAND POUNDS OF WOOD CELLULOSE OR WOOD PULP FIBER, WHICH INCLUDES A TACKIFIER, SHALL BE USED WITH HYDRAULIC SEEDING ON SLOPES 3/4:1 OR STEEPER.
4. SERICEA LESPEDEZA HAY CONTAINING MATURE SEED SHALL BE APPLIED AT A RATE OF THREE TONS PER ACRE.
5. PINE STRAW OR PINE BARK SHALL BE APPLIED AT A THICKNESS OF 3 INCHES FOR SEEDING PURPOSES. OTHER SUITABLE MATERIALS IN SUFFICIENT QUANTITY MAY BE USED WHERE ORNAMENTALS OR OTHER GROUND COVERS ARE PLANTED. THIS IS NOT APPROPRIATE FOR SEEDED AREAS.
6. WHEN USING TEMPORARY EROSION CONTROL BLANKETS OR BLOCK SOD, MULCH IS NOT REQUIRED.
7. BITUMINOUS TREATED ROVING MAY BE APPLIED ON PLANTED AREAS, ON DITCHES OR ON MATTINGS TO PREVENT EROSION. BITUMINOUS TREATED ROVING SHALL BE APPLIED WITHIN 24 HOURS AFTER AN AREA HAS BEEN PLANTED. APPLICATION RATES AND MATERIALS MUST MEET GEORGIA DEPARTMENT OF TRANSPORTATION SPECIFICATIONS.

WOOD CELLULOSE AND WOOD PULP FIBERS SHALL NOT CONTAIN GERMINATION OR GROWTH INHIBITING FACTORS. THEY SHALL BE EVENLY DISPERSED WHEN AGITATED IN WATER. THE FIBERS SHALL CONTAIN A DYE TO ALLOW VISUAL METERING AND AID IN UNIFORM APPLICATION DURING SEEDING.

APPLYING MULCH

STRAW OR HAY MULCH WILL BE SPREAD UNIFORMLY WITHIN 24 HOURS AFTER SEEDING AND/OR PLANTING. THE MULCH MAY BE SPREAD BY BLOWER–TYPE SPREADING EQUIPMENT, OTHER SPREADING EQUIPMENT OR BY HAND. MULCH SHALL BE LOOSELY PLACED TO COVER 75% OF THE SOIL SURFACE.

APPLY WOOD CELLULOSE OR WOOD FIBER MULCH SHALL BE APPLIED UNIFORMLY WITH HYDRAULIC SEEDING EQUIPMENT.

ANCHORING MULCH

ANCHOR STRAW OR HAY MULCH IMMEDIATELY AFTER APPLICATION BY ONE OF THE FOLLOWING METHODS:

1. HAY AND STRAW MULCH SHALL BE PRESSED INTO THE SOIL IMMEDIATELY AFTER THE MULCH IS SPREAD. A SPECIAL "PACKER DISK" OR DISK HARROW WITH THE DISKS SET STRAIGHT MAY BE USED. THE DISKS MAY BE SMOOTH OR SERRATED AND SHOULD BE 20 INCHES OR MORE IN DIAMETER AND 8 TO 12 INCHES APART. THE EDGES OF THE DISKS SHALL BE DULL ENOUGH TO PRESS THE MULCH INTO THE GROUND WITHOUT CUTTING IT, LEAVING MUCH OF IT IN AN ERECT POSITION. MULCH SHALL NOT BE PLOWED INTO THE SOIL.
2. SYNTHETIC TACKIFIERS, BINDERS OR HYDRAULIC MULCH SPECIFICALLY DESIGNED TO TACK STRAW, SHALL BE APPLIED IN CONJUNCTION WITH OR IMMEDIATELY AFTER THE MULCH IS SPREAD. SYNTHETIC TACKIFIERS SHALL BE MIXED AND APPLIED ACCORDING TO MANUFACTURER'S SPECIFICATIONS. ALL TACKIFIERS, BINDERS OR HYDRAULIC MULCH SPECIFICALLY DESIGNED TO TACK STRAW SHOULD BE VERIFIED AS NON–TOXIC THROUGH ITA 2021.0 TESTING. REFER TO TACKIFIERS–TAC
3. XYC OR WHEAT CAN BE INCLUDED WITH FALL AND WINTER PLANTINGS TO STABILIZE THE MULCH. THEY SHALL BE APPLIED AT A RATE OF ONE–QUARTER TO ONE–HALF BUSHEL PER ACRE.
4. PLASTIC MESH OR NETTING WITH MESH NO LARGER THAN ONE INCH BY ONE INCH MAY BE NEEDED TO ANCHOR STRAW OR HAY MULCH ON UNSTABLE SOILS AND CONCENTRATED FLOW AREAS. MATERIALS SHALL BE INSTALLED AND ANCHORED ACCORDING TO MANUFACTURER'S SPECIFICATIONS.

SEEDING MATERIAL

MULCH IS USED AS A BEDDING MATERIAL TO CONSERVE MOISTURE AND CONTROL WEEDS IN NURSERIES, ORNAMENTAL BEDS, AROUND SHRUBS, AND ON BARE AREAS ON LAWNS.

MATERIAL	DEPTH
GRAIN STRAW	4"TO 6"
GRASS HAY	4"TO 6"
PINE NEEDLES	3"TO 5"
WOOD WASTE	4"TO 6"

ABOUT 1.5 TIMES THEIR HEIGHT ARE EFFECTIVE IN CONTROLLING WIND EROSION.

CALCIUM CHLORIDE, APPLY AT RATE THAT WILL KEEP SURFACE MOIST. MAY NEED RETREATMENT.

B. PERMANENT METHODS

PERMANENT VEGETATION. SEE SPECIFI CATION DS3 –DISTURBED AREA STABILIZATION (WITH PERMANENT VEGETATION). EXISTING TREES AND LARGE SHRUBS MAY AFFORD VALUABLE PROTECTION IF LEFT IN PLACE.

TOPSOILING. THIS ENTAILS COVERING THE SURFACE WITH LESS EROSIVE SOIL MATERIAL. SEE SPECIFICATION TP – TOPSOILING.

STONE. COVER SURFACE WITH CRUSHED STONE OR COARSE GRAVEL. SEE SPECIFICATION CR–CONSTRUCTION ROAD STABILIZATION.

Ds3 DISTURBED AREA STABILIZATION (PERMANENT VEGETATION)
LOCATION IN MANUAL : 6-38

TABLE 6–5.1 FERTILIZER REQUIREMENTS

TYPES OF SPECIES	PLANTING YEAR	FERTILIZER (N–P–K)	ANALYSIS OR EQUIVALENT RATE (LBS/ACRE)	N TOP DRESSING RATE (LBS/ACRE)
1. COOL SEASON GRASSES	FIRST SECOND MAINTENANCE	6–12–12 6–12–12 10–10–10	1500 1000 400	50–100 (1)(2) – 30
2. COOL SEASON GRASSES AND LEGUMES	FIRST SECOND MAINTENANCE	6–12–12 0–10–10 0–10–10	1500 1000 400	0–50 (1) – –
3. GROUND COVERS	FIRST SECOND MAINTENANCE	6–12–12 6–12–12 10–10–10	1,300 (3) 1,300 (3) 1100	– – –
4. PINE SEEDLINGS	FIRST	20–10–5	one 21–gram pellet per seeding placed in the closing hole	–
5. SHRUB LESPEDEZA	FIRST MAINTENANCE	0–10–10 0–10–10	700 700 (4)	– –
6. TEMPORARY COVER CROPS SEEDED ALONE	FIRST	10–10–10	500	30 (5)
7. WARM SEASON GRASSES	FIRST SECOND MAINTENANCE	6–12–12 6–12–12 10–10–10	1500 800 400	50–100 (2)(6) 50–100 (2) 30
8. WARM SEASON GRASSES AND LEGUMES	FIRST SECOND MAINTENANCE	6-12-12 0-10-10 0-10-10	1500 1000 400	50 (6)

(1) APPLY IN SPRING FOLLOWING SEEDING.
(2) APPLY IN SPLIT APPLICATIONS WHEN HIGH RATES ARE USED.
(3) APPLY IN 3 SPLIT APPLICATIONS.
(4) APPLY WHEN PLANTS ARE PRUNED.
(5) APPLY TO GRASS SPECIES ONLY.
(6) APPLY WHEN PLANTS GROW TO A HEIGHT OF 2 TO 4 INCHES.

ALL AREAS TO BE SEEDED SHALL HAVE LIME APPLIED AT A RATE OF 90 LBS/1000 SF. LIME AND FERTILIZER TO BE APPLIED PRIOR TO APPLICATION OF SEED AND MIXED THOROUGHLY WITH THE SOIL.

Ds3 DISTURBED AREA STABILIZATION (PERMANENT VEGETATION)
LOCATION IN MANUAL : 6-40

TABLE 6–5.2
PLANTS, PLANTING RATES, AND PLANTING DATES FOR PERMANENT COVER OR COMPANION CROPS

SPECIES	BROADCAST per ** Acre	BROADCAST PLS * per 1000 SF	PLANTING DATES BY RESOURCE AREAS	REMARKS
BERMUDA, COMMON (Cynodon dactylon) Unhulled Seed ALONE ..... WITH OTHER PERENNIALS.....	10 lb. 6 lb.	0.2 lb. 0.1 lb.	SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	PLANT WITH WINTER ANNUALS PLANT WITH TALL FESCUE
BERMUDA, COM. (Cynodon dactylon) Hulled Seed ALONE ..... WITH OTHER PERENNIALS.....	10 lb. 6 lb.	0.2 lb. 0.1 lb.	SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	1,760,000 SEED PER POUND. QUICK COVER, LOW GROWING SOD FORMING. FULL SUN. GOOD FOR ATHLETIC FIELDS. MAY LAST FOR SEVERAL YEARS MIX W/ SERICEA LESPEDEZA
CENTIPEDE (Eremochloa ophiuroides)	SOD BLOCKS ONLY		SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	DROUGHT TOLERANT. FULL SUN OR PARTIAL SHADE. EFFECTIVE ADJACENT TO CONCRETE AND IN CONCENTRATED FLOW AREAS. IRRIGATION IS NEEDED UNTIL FULLY ESTABLISHED. DO NOT PLANT NEAR PASTURES, WHITEHEADED AS FAR AS NORTH ATHENS AND ATLANTA
FESCUE, TALL (Festuca arundinacea) ALONE ..... W/ OTHER PERENNIALS.....	50 LB 30 LB	1.1 lb. 0.7 lb.	MOUNTAINS PIEDMONT	227,000 SEED PER POUND. USE ALONE ONLY ON BETTER SITES. MIX WITH PERENNIAL LESPEDEZA OR WEEPING LOVEGRASS IN SOME OPTIONAL TALL PLANTINGS. NOT FOR HEAVY PASTURES OR ATHLETIC FIELDS
LESPEDEZA, SERICEA (Lespedeza cuneata) SCARIFIED..... UNSCARIFIED.....	60 LB 75 LB	1.4 lb. 1.7 lb.	MOUNTAINS SOUTHERN PIEDMONT SOUTHERN COSTAL PLAIN	350,000 SEEDS PER POUND. LOVEGRASS WHEN SEEDING BAHIA, OR TALL FESCUE. INOCULANT SET
LOVEGRASS, WEEPING (Eragrostis curvula) ALONE..... W/ OTHER PERENNIALS...	4 LB 2 LB	0.1 lb. 0.05 lb.	MOUNTAINS SOUTHERN